

**Fishery Conservation and Management**

**Pt. 660, Subpt. C, Table 1b**

**TABLE 1b TO PART 660, SUBPART C—2013, ALLOCATIONS BY SPECIES OR SPECIES GROUP (WEIGHTS IN METRIC TONS)**

Table 1b. To Part 660, Subpart C - 2013, Allocations by Species or Species Group. (Weights in Metric Tons)

Species	Fishery HG	Allocations			
		Trawl		Non-trawl	
		%	Mt	%	Mt
Arrowtooth flounder	4,070	95%	3,866	5%	203
Bocaccio - S of 40°10' N. lat. a/	311.6	NA	74.9	NA	236.7
Canary rockfish a/ b/	98.5	NA	52.5	NA	46.0
Chilipepper - S of 40°10 N. Lat.	1,466	75%	1,100	25%	367
Cowcod - S of 40°10' N. lat. a/	2.9	NA	1.0	NA	1.9
Darkblotched rockfish c/	296.2	95%	281.4	5%	14.8
Dover sole	23,410	95%	22,240	5%	1,171
English sole	6,712	95%	6,376	5%	336
Lingcod					
N of 40°10' N. lat.	2,758	45%	1,241	55%	1,517
S of 40°10' N. lat.	1,102	45%	496	55%	606
Longnose Skate a/	1,928	90%	1,735	10%	193
Longspine thornyhead					
N of 34°27' N. lat.	1,963	95%	1,865	5%	98
Minor shelf rockfish north a/	903	60.2%	543	39.8%	359
Minor shelf rockfish south a/	668	12.2%	81	87.8%	587
Minor slope rockfish north	1,098	81%	889	19%	209
Minor slope rockfish south	597	63%	376	37%	221
Other flatfish	4,682	90%	4,214	10%	468
Pacific cod	1,191	95%	1,131	5%	60
POP - N of 40°10' N. lat. d/	133.5	95%	126.8	5%	6.7
Pacific whiting	TBA	100%	TBA	0%	TBA
Petrale sole a/	2,358.0	NA	2323.0	NA	35.0
Sablefish					
N of 36° N. lat.		See Table 1c of this subpart			
S of 36° N. lat.	1,434	42%	602	58%	832
Shortspine thornyhead					
N of 34°27' N. lat.	1,481	95%	1,407	5%	74
S of 34°27' N. lat.	355	NA	50	NA	305
Splitnose - S of 40°10 N. Lat.	1,598	95%	1,518	5%	80
Starry Flounder	1,513	50%	757	50%	757
Widow e/	1,411	91%	1,284	9%	127
Yelloweye rockfish a/	12.2	NA	1.0	NA	11.2
Yellowtail - N of 40°10 N. Lat.	3,677	88%	3,235	12%	441

a/ Allocations decided through the biennial specification process.

b/ 12.6 mt of the total trawl allocation of canary rockfish is allocated to the at-sea whiting fisheries, as follows: 5.2 mt for the mothership fishery, and 7.4 mt for the catcher/processor fishery.

c/ 9 percent (25.3 mt) of the total trawl allocation for darkblotched rockfish is allocated to the whiting fisheries, as follows: 10.6 mt for the shorebased IFQ fishery, 6.1 mt for the mothership fishery, and 8.6 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at § 660.140(d)(1)(ii)(D).

d/ 30 mt of the total trawl allocation for POP is allocated to the whiting fisheries, as follows: 12.6 mt for the shorebased IFQ fishery, 7.2 mt for the mothership fishery, and 10.2 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at § 660.140(d)(1)(ii)(D).

e/ 500 mt of the total trawl allocation for widow rockfish is allocated to the whiting fisheries, as follows: 210 mt for the shorebased IFQ fishery, 120 mt for the mothership fishery, and 170 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at § 660.140(d)(1)(ii)(D).

[78 FR 592, Jan. 3, 2013]